


# *Astrophysical Neutrinos and New Physics with Water Cherenkov detectors*





Kenny, Chun Yu Ng (吳震宇)  
Weizmann Institute of Science



# Neutrinos Astro-particle physics

## Astro ----- Particle




- Reveal concealed astrophysical sites
  - Solar, Supernova 
  - AGN
- Ideal messenger
- Cosmic hadron accelerator
  - 100 year CR problem
- Neutrino mass 
  - Dirac mass, why so small?  
-> Sterile neutrino!
  - Majorana mass?
  - Both?
- New physics portals?
  - Dark matter
  - Secret interactions

# Supernova Neutrinos



# Galactic Supernova

- SN1987A



Vissani 2015


Blum Kushnir 2016



# Galactic Supernova



Tamborra+2013

- IceCube




Halzen Raffelt 2009

- Bounce time  $\pm 3.5\text{ms}$   $\rightarrow$  GW



# Nearby Supernova

- Can we see nearby supernova?
  - Local Overdensity
- 1/year <6Mpc
- Singles (+ optical)
  - 30%
- 10 years
  - $1 - P(\text{no det.}) > 97\%$




Nakamura et al 2016

# Diffuse supernova neutrino background

## Average neutrino emission

- Use >100 simulations to characterize progenitor dependence of neutrinos
- Include collapse to black holes, characterized by critical compactness




reviews by Beacom (2010), Lunardini (2010)

Shunsaku Horiuchi (VT CNP)

## Event rate predictions


Hyper-K sensitive to small compactness ( $\xi_{2.5} < 0.2$ , or  $f_{BH} > 0.2$ )

Spectrum	SK + Gd (>10MeV) [/yr]
4 MeV	1.8 +/- 0.5
4 MeV+BH	< 2.5
SN1987A	1.7 +/- 0.5



Horiuchi et al (2018); see also Lunardini (2009), Lien et al (2010), Moller et al (2018)


# Astrophysical Neutrinos




# Diffuse Astrophysical Neutrinos

Schneider TeVPA2018

- IceCube HESE and  $\nu_\mu$
- $E^{-2.2}$  vs  $E^{-2.9}$  ?
- Two components?




## Differential Flux






- Source cannot be rare and bright
- Or maybe hidden?

# Multi-messenger Connection




Ahlers, Halzen 2018

# Where are the Taus?



- Double Bang
- Double Pulse





IceCube 2015  
3 years



# Point source astrophysical Neutrinos

- Era of multimessenger astronomy with TXS events




Blazar as a hadronic accelerator!

Association  $\sim 3$  sigma

Neutrino flare  $\sim 3.5$  sigma

Need more of these!







# Sun – Cosmic-ray beam dump


Seckel, Stanev, Gaisser (1991),  
Moskalenko, Karakula (1993),  
Ingelman, Thunman (1996), +


Low density atmosphere  
-> long interaction length  
-> more decays  
-> Higher flux, higher energy



# Neutrino Flux


- Showers:  $\nu_{e,\tau}CC, \nu_X NC$
- Tracks:  $\nu_\mu CC$
- $\nu_\mu CC$  for directionality
  - kinematic angle
- Above  $\sim 3$  TeV, greater than Earth ATM background
- Unclear how solar magnetic fields change the prediction






# $SA\nu$ as a Signal

- Muon ( $>1\text{TeV}$ )
  - Energy resolution via energy loss
- $\sim 5$  signal events in 10 years (4 bkg)
- ~~Another~~ 1<sup>st</sup> high-energy neutrino source?
- Common source for IceCube + KM3NeT



# Solar Atmospheric gamma rays



Neutrinos could help understand the gamma rays

Tang et al 2018


- Time variation
- Hard spectrum
- Large flux

# Dark Matter and New physics




# Dark Matter

- Weakly Interacting Massive Particles (WIMPs)




- Abundance  $\leftrightarrow$  **Total** Cross section!

$$\frac{\Omega_{DM} h^2}{0.1} \approx \frac{3 \times 10^{-26} \text{ cm}^3 \text{ s}^{-1}}{\langle \sigma v \rangle}$$




# Dark Matter Annihilation

Antiproton, Reinert & Winkler 2018



Gamma rays, Fermi collab. 2017




- Specific channels or models




# The simplest WIMP hypothesis

- Total cross section constraint
  - Arbitrary, mixed channel (mixed spectrum)
  - Fermi dwarf, AMS positron, Planck CMB
- New physics for large  $\sigma v$
- Sub DM thermal relics
- All visible channels except **Neutrinos!**


Leane, Slatyer, Beacom, KCYN, 2018



# Neutrino Channel



Klop Ando 1809.00671  
 Also see  
 Olivares-Del Campo, Palomares-Ruiz, Pascoli 1805.09830





Mijakowski TAUP 2017

- Reaching thermal?
  - A significant milestone for testing WIMPs


# Dark Matter

- Weakly Interacting Massive Particles

$\chi$





$\nu$




# Dark Matter Search from the Sun

Rott, NOW 2018




# Solar ATM neutrino – indirect detection Neutrino Floor



No B-field effect are considered

IceCube Search ongoing [S. In & C. Rott ICRC17 (965)]



KCYN, Beacom, Peter, Rott, 1703.10280

See also



Arguelles+ 1703.07798

Edsjo+ 1704.02892

# Dark Matter with long-lived mediators


Leane, KCYN, Beacom 1703.04629

- No neutrino absorption
- + EM signatures!





# Dark Matter Decay

Cohen, Murase, Rodd, Safdi, Soreq 2017




IceCube Collab, 2018




Chianese, Miele, Morisi 2017

# IceCube Boosted Dark Matter


Following search proposed by [Kopp, Liu, Wan \(2015\)](#)  
using “Echo Technique” [Li, Bustamante, Beacom \(2016\)](#)




Very heavy dark matter particle  $\phi$  decays to lighter stable dark matter  $\chi \rightarrow \text{boost!}$



Recoil  
(only hadronic  
cascades)



$\phi \rightarrow \chi \bar{\chi}$ ,  $a \rightarrow b \bar{b} \rightarrow \nu's$



Neutrons capture on hydrogen and product 2.2 MeV gamma. In seawater, 33% of neutrons capture on Cl; the emitted gamma rays have 8.6 MeV, making the neutron echoes more visible


“Echo Technique” holds prospects to individually tag high-energy NC and CC interactions !

*May sound crazy, but is just an example for exotic interactions in IceCube detectable via recoil*


see also [A. Steuer, L. Koepke \[IceCube\] PoS\(ICRC2017\)1008](#)




# Super-K Boosted Dark Matter




161.9 kiloton-years of Super-Kamiokande IV data



Cone search: 8 cones from 5° to 40° around GC  
→ no cluster found around Galactic Center






90% limits for  
 $m_{\gamma'} = 20 \text{ MeV}$   
 $m_B = 200 \text{ MeV}$   
 $g' = 0.5$





• < 1.33 GeV + 1.33 GeV-20 GeV • > 20 GeV

# Cosmic neutrino cascades

- Secret neutrino interactions






# Neutrino Dark Matter Interaction



Argüelles, Kheirandish, Vincent, 2017

# Dark Matter Beam (T2K)




deNiverville et al, 2017

# Summary

- Rich astroparticle physics, many can only be done with water Cherenkov detectors
- Exciting times ahead for new detectors and maybe new techniques


Thanks!

# Backup slides



# Solar Atmospheric Neutrino Floor

- Large direct detection experiments are needed to reach  $10^{-44} \text{ cm}^2$



Solar, diffuse supernova, atmospheric neutrinos  
Ruppin et al. 2014

