

## The FELBE THz/IR FEL : Overview of the Facility and User Activities

The FELBE User Facility at the ELBE Center for High-Power Radiation Sources offers a pair of FELs that deliver beam to eight different user labs. The FELs are driven by a two-stage Superconducting RF (SRF) linac, which produces a quasi-CW beam (13 MHz/1 mA) at an energy of up to 36 MeV. The tuning range spanned by the two FELs extends from the mid IR to THz (5 – 250  $\mu\text{m}$ ). The spectral range and ultrashort pulse width ( $\tau_p \approx 0.7 - 25$  ps) are ideal for time-resolved measurements of many types of transient processes in low-dimensional materials [1], quantum structures [2], and correlated systems [3]. The high pulse energy can also drive nonlinear phenomena [4] and strong coupling [5] in light-matter interactions. The FELBE User Labs are equipped with instrumentation and synchronized ultrashort table-top lasers (i.e. Ti:Sa oscillators, regens, OPAs, SFG/DFG) which facilitate various classes of degenerate (single-color), and non-degenerate (two-color) pump-probe experiments. Optical cryostats and an 8 T split coil magnet are also available for low temperature and magnetic field dependent studies. Furthermore, the FELBE beamline extends into the adjacent High Field Magnet Lab (HLD) for performing magneto-optical spectroscopy measurements at fields up to 70 T [6]. The high repetition rate and tunability of the FELBE beam has uniquely enabled revolutionary methods in scattering-Scanning Nearfield Optical Microscopy (s-SNOM) to image novel light-matter interactions with resolution far below the diffraction limit [7]. Proposals for beamtime on FELBE and the other secondary sources at ELBE are invited from users twice a year.

(<https://www.hzdr.de/FELBE>).

- [1] T. Venanzi, et al., ACS Photonics **8**, 2931-2939 (2021).
- [2] J. Schmidt, et al., Optics Express **28**, 25358-25370 (2020).
- [3] M. M. Jadidi, et al., Phys. Rev. B **102**, 245123 (2020).
- [4] F. Meng, et al., Phys. Rev. B **102**, 075205 (2020).
- [5] B. Piętko, et al., Phys. Rev. Lett. **119**, 077403 (2017).
- [6] M. Ozerov, et al., Phys. Rev. Lett. **113**, 157205 (2014).
- [7] T. V. A. G. de Oliveira, et al., Adv. Mater. **33**, 2005777 (2021).

**Primary author(s) :** Dr KLOPF, J. Michael (Helmholtz-Zentrum Dresden-Rossendorf (HZDR))

**Co-author(s) :** Dr EVTUSHENKO, Pavel (Helmholtz-Zentrum Dresden-Rossendorf (HZDR)); Prof. HELM, Manfred (Helmholtz-Zentrum Dresden-Rossendorf (HZDR)); Dr KEHR, Susanne C. (Technische Universität Dresden); Dr LEHNERT, Ulf (Helmholtz-Zentrum Dresden-Rossendorf (HZDR)); Prof. MICHEL, Peter (Helmholtz-Zentrum Dresden-Rossendorf (HZDR)); Dr PASHKIN, Alexej (Helmholtz-Zentrum Dresden-Rossendorf (HZDR)); Dr WINNERL, Stephan (Helmholtz-Zentrum Dresden-Rossendorf (HZDR)); Dr ZVYAGIN, Sergei (Helmholtz-Zentrum Dresden-Rossendorf (HZDR))

**Presenter(s) :** Dr KLOPF, J. Michael (Helmholtz-Zentrum Dresden-Rossendorf (HZDR))

**Session Classification :** Tue 05/07 Morning/ Change Abstract ID