

## **The back – gated AlGa<sub>N</sub>/Ga<sub>N</sub> HEMT structure for observation of twisted plasmonic states in terahertz frequency range.**

*Wednesday, 6 July 2022 17:35 (15)*

The Inverse Faraday Effect (IFE) is the appearance of stationary magnetic moments magnetization caused by circularly polarized light. Up to this moment, IFE has been mostly studied in the magnetic materials. In recent years, the IFE was predicted in the periodic lattice of metallic disks or spheres placed in the vicinity of two-dimensional electron liquid and under illumination of the external circularly polarized light. The radiation causes the DC current loops in the electron liquid, thus leading to appearance of static magnetic moments. As a result, the interaction between metal disks and two-dimensional electron liquid, the “twisted” plasmonic modes are excited what leads to the appearance of DC circulating current due to rectification. In this work, we present the basic idea of IFE. The mechanism of the effect is described and the theoretical predictions are presented. We propose Ga<sub>N</sub>/AlGa<sub>N</sub> as a basic system for the experimental realization of the IFE. We present Ga<sub>N</sub>/AlGa<sub>N</sub> HEMT like structure with 2-dimensional electron gas as a channel, that was made in order to observe the twisted plasmonic modes experimentally. On top of the structure, the periodic lattice of metal disks was fabricated with use of electron beam lithography. We present the theoretical predictions and the technological realization of the structure as well as its basic characterization. The frequency of the twisted plasmonic modes is expected to be in range of 0.6 - 1.2 THz thus leading to potential applications in terahertz physics and technology.

**Primary author(s) :** Dr DRÓŹDŹ, Piotr (CENTERA Laboratories, Institute of High Pressure Physics, PAS)

**Co-author(s) :** Prof. KACHOROVSKII, Valentin (CENTERA Laboratories, Institute of High Pressure Physics PAS, Warsaw, Poland; Ioffe Institute, St. Petersburg, Russia; Rensselaer Polytechnic Institute, Troy, New York, USA); Dr PRYSTAWKO, Pawel (Institute of High Pressure Physics); Mr SŁOWIKOWSKI, M. (CEZAMAT, Warsaw University of Technology, 02-822 Warsaw, Poland); Mr FILIPIAK, M. (CEZAMAT, Warsaw University of Technology, 02-822 Warsaw, Poland); Dr YAVORSKI, Dymitri (CENTERA Laboratories, Institute of High Pressure Physics PAS, Warsaw, Poland; Faculty of Physics, University of Warsaw, Warsaw, Poland); Mrs SZOŁA, Maria (CENTERA Laboratories, Institute of High Pressure Physics PAS, Warsaw, Poland; Laboratoire Charles Coulomb (L2C), UMR 5221 CNRS-Université de Montpellier, F-34095 Montpellier, France); Prof. KNAP, Wojciech (CENTERA Laboratories, Institute of High Pressure Physics, PAS)

**Presenter(s) :** Dr DRÓŹDŹ, Piotr (CENTERA Laboratories, Institute of High Pressure Physics, PAS)

**Session Classification :** Wed 06/07 Poster Session/ ID: