

# **Particle Astrophysics in Poland**

Monday 20 May 2019 - Tuesday 21 May 2019

## **Book of Abstracts**



# Contents

A signature of sterile neutrino dark matter in the Local Universe . . . . .	1
ATHENA - upgrade . . . . .	1
AstroCeNT . . . . .	1
Bayesian reconstruction of the Milky Way dark matter distribution . . . . .	1
Bifurcations in Ratra-Peebles quintessence models and their extensions . . . . .	1
CREDO . . . . .	1
CTA . . . . .	1
Comprehensive measurement of pp-chain solar neutrinos in Borexino . . . . .	1
Constraining cosmological parameters using reverberation-measured quasars . . . . .	2
Constraining the halo size from possible density profiles of hydrogen gas of Milky Way Galaxy . . . . .	2
Cosmic Ray shower simulations for KM3NeT . . . . .	2
Cosmic-ray ensembles: from formation to monitoring . . . . .	2
Dark matter haloes and their galaxies at $z\sim 3$ . . . . .	2
Dark matter searches at Super-Kamiokande and its extensions . . . . .	3
EGO and European Road Map for Astroparticle Physics . . . . .	3
Electroweak baryogenesis and dark matter from a complex singlet scalar . . . . .	3
Environmental dependence of properties of galaxies in Galaxy and Mass Assembly (GAMA) survey and its evolution using Marked statistics . . . . .	3
HESS . . . . .	4
Heliospheric modulation of cosmic rays . . . . .	4
Higgs domain walls in the thermal background . . . . .	4
Instabilities and nonlinear interactions of scalar fields perturbations in models of cosmological inflation . . . . .	4
International Research Agenda and other programs of FNP . . . . .	4

JEM-EUSO . . . . .	4
KM3NeT: scientific prospects and current status . . . . .	4
Ligo/Virgo observational results . . . . .	4
MAGIC, CTA LST . . . . .	4
MeV Space Observatories . . . . .	5
Meeting of KRAC (closed) . . . . .	5
Missing black holes and the dark matter puzzle . . . . .	5
Official remarks . . . . .	5
Opening . . . . .	5
POLAR - detector for Gamma Ray bursts polarization . . . . .	5
Pierre Auger Observatory . . . . .	5
Production of Purely Gravitational Vector Dark Matter . . . . .	5
Rotating neutron stars as sources of gravitational waves . . . . .	6
Search for Axionlike Dark Matter through Nuclear Spin Precession in Electric and Magnetic Fields . . . . .	6
Searches for the Majorana neutrino in GERDA . . . . .	6
Searching for continuous gravitational wave signals from rotating neutron stars . . . . .	6
Searching for long-lived particles from light dark sectors . . . . .	6
T2K . . . . .	6
Testing dark matter with Cherenkov light – prospects of H.E.S.S. and CTA for exploring minimal supersymmetry . . . . .	7
Testing dark matter with Cherenkov light – prospects of H.E.S.S. and CTA for exploring minimal supersymmetry . . . . .	7
The DarkSide Program of Dark Matter Searches and Its Broader Impact . . . . .	7
The origin of the coalescing compact object binaries . . . . .	7
Theory of Acceleration of Particles . . . . .	7
Theory of Cosmic Ray Origin . . . . .	7
Theory of GRBs . . . . .	7
Theory of Pulsars and Pulsar Wind Nebulae . . . . .	7
Theory of high energy processes in Binary Systems . . . . .	8
Time variability of low angular momentum accretion flows around black hole. . . . .	8

Toward precise and accurate determination of the Hubble constant based on supernovae Ia, Cefeids and eclipsing binaries . . . . .	8
Understanding Our Universe from Deep Underground . . . . .	8
Welcome . . . . .	9
eXTP X-ray time and polarymetry mission . . . . .	9



**Dark Matter and Dark Energy in Astronomical Measurements / 40**

## **A signature of sterile neutrino dark matter in the Local Universe**

**High Energy Astrophysics from Space / 34**

## **ATHENA - upgrade**

**Dark Matter, Astrocent, KRAC / 61**

## **AstroCeNT**

**Theory (Dark Matter, Dark Energy, Early Universe) / 49**

## **Bayesian reconstruction of the Milky Way dark matter distribution**

**Theory (Dark Matter, Dark Energy, Early Universe) / 50**

## **Bifurcations in Ratra-Peebles quintessence models and their extensions**

**Cosmic Rays / 26**

## **CREDO**

**Gamma Ray Astronomy / 21**

## **CTA**

**Neutrino Physics / 14**

## **Comprehensive measurement of pp-chain solar neutrinos in Borexino**

**Dark Matter and Dark Energy in Astronomical Measurements / 42****Constraining cosmological parameters using reverberation-measured quasars**

5

**Constraining the halo size from possible density profiles of hydrogen gas of Milky Way Galaxy**Sayan Biswas<sup>1</sup> ; Nayantara Gupta<sup>1</sup><sup>1</sup> *Raman Research Institute*

Galactic magnetic field (GMF) and secondary cosmic rays (CRs) (e.g. <sup>10</sup>beryllium, boron, antiproton) are important components to understand the propagation of CRs in the Milky Way Galaxy. Realistic modeling of GMF is based on the Faraday rotation measurements of various Galactic and extragalactic radio sources and synchrotron emission from CR leptons in the radio frequency range, thereby providing information of halo height. On the other hand, diffusion coefficient and halo height are also estimated from the <sup>10</sup>Be/<sup>9</sup>Be and B/C ratios. Moreover, density distribution of gaseous components of interstellar medium (ISM) also plays an important role as secondary CRs are produced due to interaction of primary CRs with the gaseous components of ISM. We consider mainly molecular, atomic, and ionized components of hydrogen gas for our study. Recent observations and hydrodynamical simulations provide new forms of density profiles of hydrogen gas in Milky Way Galaxy. In the `DRAGON` code, we have implemented our chosen density profiles, based on realistic observations in radio, X-ray and  $\gamma$ -ray wavebands, and hydrodynamical simulations of interstellar hydrogen gas to study the variation in the height of the halo required to fit the observed CR spectra. Our results show the halo height ( $z_t$ ) varies in the range of 2 to 6 kpc for the density profiles considered in our work.

**Session:**

Cosmic Rays

**Neutrino Physics / 18****Cosmic Ray shower simulations for KM3NeT**

52

**Cosmic-ray ensembles: from formation to monitoring****Session:****Dark Matter and Dark Energy in Astronomical Measurements / 43****Dark matter haloes and their galaxies at  $z \sim 3$**



**Neutrino Physics / 16****Dark matter searches at Super-Kamiokande and its extensions****Dark Matter, Astrocent, KRAC / 59****EGO and European Road Map for Astroparticle Physics****Theory (Dark Matter, Dark Energy, Early Universe) / 45****Electroweak baryogenesis and dark matter from a complex singlet scalar**

8

**Environmental dependence of properties of galaxies in Galaxy and Mass Assembly (GAMA) survey and its evolution using Marked statistics**Unnikrishnan Sureshkumar<sup>1</sup><sup>1</sup> *Astronomical Observatory of the Jagiellonian University, Krakow***Corresponding Author(s):** [usureshkumar@oa.uj.edu.pl](mailto:usureshkumar@oa.uj.edu.pl)

Observed galaxies trace an underlying network of gravitationally dominant dark matter; we know however that they trace it in a biased way, and that this bias depends on galaxy properties. In many studies galaxy luminosity and/or galaxy stellar mass is used as a convenient (even if also biased) proxy of its host dark matter halo; in the same time it was also observed that clustering of luminosity-selected and stellar mass-selected samples is not identical, especially at higher redshifts (see Marulli et al. 2013; Durkalec et al. 2018), and these differences are quite complex. We use marked correlation function as a tool to study small-scale galaxy clustering weighted by these two properties. We present our first results from the study of the dependence of galaxy clustering on luminosity and stellar mass in the redshift range  $0.1 < z < 0.5$  using 54262 galaxies from the Galaxy And Mass Assembly (GAMA) survey, covering a total area of 180 sq. deg. We measure the real space luminosity-marked and stellar mass-marked correlation functions for a set of volume-limited subsamples selected by the absolute magnitude and stellar mass. We present the results of a comparative study of both the properties with the aim to show how these two properties trace dark matter halo mass and local density field.

**Session:**

Dark Matter and Dark Energy in Astronomical Measurements

**Gamma Ray Astronomy / 20**

## **HESS**

**Cosmic Rays / 23**

## **Heliospheric modulation of cosmic rays**

**Theory (Dark Matter, Dark Energy, Early Universe) / 46**

## **Higgs domain walls in the thermal background**

**Theory (Dark Matter, Dark Energy, Early Universe) / 44**

## **Instabilities and nonlinear interactions of scalar fields perturbations in models of cosmological inflation**

**Dark Matter, Astrocent, KRAC / 57**

## **International Research Agenda and other programs of FNP**

**Cosmic Rays / 25**

## **JEM-EUSO**

**Neutrino Physics / 17**

## **KM3NeT: scientific prospects and current status**

**Gravitational Waves / 35**

## **Ligo/Virgo observational results**

**Gamma Ray Astronomy / 22**

**MAGIC, CTA LST**

**High Energy Astrophysics from Space / 33**

**MeV Space Observatories**

2

**Meeting of KRAC (closed)**

**Dark Matter and Dark Energy in Astronomical Measurements / 39**

**Missing black holes and the dark matter puzzle**

**Dark Matter, Astrocent, KRAC / 56**

**Official remarks**

1

**Opening**

**High Energy Astrophysics from Space / 54**

**POLAR - detector for Gamma Ray bursts polarization**

**Cosmic Rays / 24**

**Pierre Auger Observatory**

51

## **Production of Purely Gravitational Vector Dark Matter**

**Session:**

**Gravitational Waves / 38**

### **Rotating neutron stars as sources of gravitational waves**

62

### **Search for Axionlike Dark Matter through Nuclear Spin Precession in Electric and Magnetic Fields**

Data collected in the measurement of the neutron electric dipole moment (nEDM) were used to look for the ultralow-mass axionlike dark matter. Analysis of ratio of frequencies of spin-precession of ultracold neutrons and  $^{199}\text{Hg}$  atoms enabled us to estimate an axion-induced oscillating electric dipole moment of the neutron and an axion-wind spin-precession effect. Our null result sets the first laboratory constraints on the coupling of axion dark matter to gluons, which improve on astrophysical limits by up to 3 orders of magnitude, and also improves on previous laboratory constraints on the axion coupling to nucleons by up to a factor of 40.

**Session:**

**Neutrino Physics / 15**

### **Searches for the Majorana neutrino in GERDA**

**Gravitational Waves / 37**

### **Searching for continuous gravitational wave signals from rotating neutron stars**

**Theory (Dark Matter, Dark Energy, Early Universe) / 48**

### **Searching for long-lived particles from light dark sectors**

**Neutrino Physics / 19**

**T2K**

53

## **Testing dark matter with Cherenkov light – prospects of H.E.S.S. and CTA for exploring minimal supersymmetry**

**Co-author(s):** A. Hryczuk ; E. Moulin ; E.M. Sessolo ; K. Jodłowski ; L. Rinchiuso ; L. Roszkowski ; S. Trojanowski

**Session:**

Theory (Dark Matter, Dark Energy, Early Universe) / 47

## **Testing dark matter with Cherenkov light – prospects of H.E.S.S. and CTA for exploring minimal supersymmetry**

Dark Matter, Astrocent, KRAC / 60

## **The DarkSide Program of Dark Matter Searches and Its Broader Impact**

Gravitational Waves / 36

## **The origin of the coalescing compact object binaries**

High Energy Astrophysics / 29

## **Theory of Acceleration of Particles**

High Energy Astrophysics / 30

## **Theory of Cosmic Ray Origin**

High Energy Astrophysics / 31

## **Theory of GRBs**

High Energy Astrophysics / 28

## Theory of Pulsars and Pulsar Wind Nebulae

High Energy Astrophysics / 27

### Theory of high energy processes in Binary Systems

6

### Time variability of low angular momentum accretion flows around black hole.

Ishika Palit<sup>1</sup> ; Agnieszka Janiuk<sup>1</sup> ; Petra Sukova<sup>2</sup>

<sup>1</sup> *Center for Theoretical Physics of Polish Academy of science*

<sup>2</sup> *Astronomical Institute, Czech Academy of Sciences*

**Corresponding Author(s):** ishi2694@gmail.com, pet.mff@gmail.com, agnes@cft.edu.pl

Observations show the emission of hard and soft photons at high energies (X-ray or gamma ray) in the black hole accretion flow's spectra. These hard photons are observed at very high frequency which implies that they are produced near black hole horizon. As the quality and quantity of the high energy observations improved over the years, evidence mounted showing that photons must be created in a hot, tenuous, advection dominated region called the corona. This corona, boiling violently above the comparatively cool disk, is very close to the event horizon of the black hole. A relativistic fluid flowing into the black hole must have a varying adiabatic index rather than a constant one throughout the accretion disk.

Our recent work present the relativistic 2D simulation of such axisymmetric, inviscid, hydrodynamic accretion flows in a fixed Kerr black hole gravitational field. The flow is considered to have low angular momentum with respect to Keplerian one. In quasi-spherical, transonic accretion flow, occurrence and location of shock and sonic points depends on the parameters of the flow. Studying the evolution of this kind of flow with time shows oscillation of shock position in response to pressure against rotational force for some particular parameter space. I will talk about such oscillatory behavior of shock position and respective effect on mass accretion rate and frequency with varying adiabatic index. I will also discuss the relevance of our results with the observed phenomenon - QPO's (Quasi periodic oscillations) from galactic black holes and micro-quasars

**Session:**

High Energy Astrophysics

Dark Matter and Dark Energy in Astronomical Measurements / 41

### Toward precise and accurate determination of the Hubble constant based on supernovae Ia, Cefeids and eclipsing binaries

Dark Matter, Astrocent, KRAC / 58

### Understanding Our Universe from Deep Underground

**Dark Matter, Astrocent, KRAC / 55**

**Welcome**

**High Energy Astrophysics from Space / 32**

**eXTP X-ray time and polarymetry mission**